

DOI: <https://doi.org/10.17816/DD79504>

Osteopoikilosis in the Ribs, Pelvic Region and Spine: A Case Report

Paparella MT¹, Gangai I¹, Porro C¹, Eusebi L², Silveri F³, Cammarota A⁴, Guglielmi G^{4,5}

AFFILIATIONS

¹Department of Clinical and Experimental Medicine, Foggia University School of Medicine, Viale L. Pinto 1, 71121 Foggia, Italy

²Radiology Unit, "Carlo Urbani"Jesi, Italy

³Rheumatology Unit, University of Ancona, Italy

⁴IRCCS-CROB, Rionero in Vulture, Italy

⁵Radiology Unit, Barletta University Hospital, Italy

ABSTRACT

Osteopoikilosis is a rare inherited benign bone dysplasia incidentally found on radiological exams. It is characterized by a specific radiological pattern: diffuse, round or oval, symmetrically shaped sclerotic bone areas distributed throughout the skeleton. It is essential to do a correct diagnosis because these lesions could be easily confused with bone metastasis. We reported a case of an osteopoikilosis patient presenting to our clinic with transient loss of consciousness and without any numbness, tingling and weakness in the legs or other parts of the body. The computed tomography scan showed multiple small sclerotic foci bone islands, scattered throughout the thoracic and lumbar spine, ribs, pelvic bone, sacrum and bilateral proximal femur. No significant increase in the activity was detected in technetium-99m whole-body bone scintigraphy. The patient was diagnosed with characteristic radiological findings of osteopoikilosis and was followed up.

Keywords: osteopoikilosis, bone dysplasia, clinical case.

To cite this article

Paparella MT, Gangai I, Porro C, Eusebi L, Silveri F, Cammarota A, Guglielmi G. Osteopoikilosis in the Ribs, Pelvic Region and Spine: A Case Report. Digital Diagnostics. 2021;2(4):00-00. DOI: <https://doi.org/10.17816/DD79504>

Received: 03.09.2021

Accepted: 16.11.2021

Published: 07.12.2021

BACKGROUND

Osteopoikilosis is a rare benign bone dysplasia that affects about one in every 50,000 people, usually with no age or gender differences [1].

It is characterized by numerous circular or ovoid sclerotic bone lesions symmetrically distributed throughout the skeleton [2]. Lesions are frequently found incidentally on imaging studies for unrelated complaints [3]. Histologically, the lesions are thicker trabeculae of lamellar osseous tissue with haversian systems within the cancellous structure; they are most likely foci of bone that did not become cancellous throughout growth and differentiation. The condensation of cancellous bone in osteopoikilosis consists of a peripheral area of trabeculae in which osteocytes are scant, and there

are no osteoblasts or osteoclasts (both are present in the central core of irregular trabeculae) [4,5]. We report a case of osteopoikilosis patient who presented to our clinic for a syncope.

DESCRIPTION OF THE CASE

A 43-year-old female patient was taken to the emergency room by ambulance after experiencing transient loss of consciousness. The initial evaluation consisting of history, physical examination, 12-lead electrocardiogram and laboratory tests did not reveal any abnormalities; thus, a total-body computed tomography (CT) was performed. The CT scan showed multiple small sclerotic foci bone islands, scattered throughout the thoracic (Figure 1a) and lumbar spine (Fig. 1b), ribs, pelvic bone (Fig. 2), sacrum (Fig. 3) and bilateral proximal femur (Fig. 4). All bones were free of any cortical erosion or periosteal reaction. No other signs, such as rubor or edema, were noticed; moreover, the patient did not describe any numbness, tingling and weakness in the legs or other parts of the body. The CT pattern was suspicious for osteopoikilosis. The relative clinical and laboratory tests, such as routine blood count, erythrocyte sedimentation rate, serum electrolytes, tumor markers, alkaline and acid phosphatase, ANA and anti-DS-DNA were negative for any type of arthritis, infection or osteoblastic bone metastases, which were in the differential diagnosis. No significant increase in the activity was detected in technetium-99m whole-body bone scintigraphy. The patient was diagnosed with typical radiological findings of osteopoikilosis by excluding other differential diagnoses and was followed up.

DISCUSSION

Osteopoikilosis (also known as “spotted bone disease” or osteopathia condensans disseminata) is a rare bone dysplasia and was first described by Albers-Schönberg in 1915 [6]. The incidence of this disease is estimated around one in 50,000, usually without age or gender differences [1]. It is usually autosomal dominant in inheritance, but sporadic forms are also reported [1]. Current literatures suggest loss-of-function mutations of *LEM domain-containing 3 (LEMD3)* gene located on 12q might be the cause. These mutations could also affect soft tissue and skin, causing melorheostosis that is a benign sclerosing bone dysplasia with cortical hyperostosis, thickening and fibrosis of overlying skin and Buschke–Ollendorff syndrome that comprises osteopoikilosis associated with disseminated connective tissue and cutaneous yellowish nevi [7,8]. Osteopoikilosis lesions are typically found incidentally on imaging studies done for unrelated complaints [3]. Radiological lesions of osteopoikilosis are typical: they are characterized by numerous symmetrical, homogeneous, well circumscribed, small (1–10 mm in diameter) and round or oval shaped sclerotic lesions. The most commonly affected areas are the epiphyses of short tubular bones and the metaphyses of long bones. In addition, carpal and tarsal bones, scapula, pelvis and sacrum are reported to be frequently affected [9,10]. Ribs, clavicles, spine and skull involvement is uncommon [11]. Because of their similarities, the radiological lesions of osteopoikilosis can be confused with osteoblastic bone metastases, but there are significant differences that allow us to make a differential diagnosis. In contrast to bone metastasis, the sclerotic lesions in osteopoikilosis are symmetrical, consistent in size and do not induce cortical erosion. As a result, bone scintigraphy plays an important role in definitive diagnosis; in fact, a normal radionuclide bone scan generally excludes the possibility of osteoblastic bone metastasis. Nevertheless, several cases of osteopoikilosis with an abnormal bone scan have been reported in the literature [12,13].

CONCLUSION

Although osteopoikilosis is a rare condition, it can be easily diagnosed through its typical radiological findings. Therefore, clinicians must be aware of and recognize this image pattern in order to make an accurate diagnosis and prevent further examinations and aggressive treatments.

ADDITIONAL INFORMATION


Funding source. This study was not supported by any external sources of funding

Competing interests. The authors declare that they have no competing interests.

Authors' contribution. All authors made a substantial contribution to the conception of the work, acquisition, analysis, interpretation of data for the work, drafting and revising the work, final approval of the version to be published and agree to be accountable for all aspects of the work. Paparella MT and Gangai I - contributed equally to the research work related to the topic and the manuscript writing; Porro C, Eusebi L and Silveri F - literature research and data acquisition; Cammarota A and Guglielmi G - critical revision of manuscript.


Consent for publication. Written consent was obtained from the patient for publication of relevant medical information and all of accompanying images within the manuscript.

FIGURES



1a

Fig. 1a. Transverse cross-section computed tomography scan passing through the thoracic (A) and lumbar (B) spine. It shows numerous, well-defined, homogeneous, circular, hyperdense foci in spinous processes and vertebral arches.



1b

Fig. 1b. Transverse cross-section computed tomography scan passing through the thoracic (A) and lumbar (B) spine. It shows numerous, well-defined, homogeneous, circular, hyperdense foci in spinous processes and vertebral arches.




Fig. 2. Transverse cross-section computed tomography scan passing through the seventh rib. It shows numerous hyperdense lesions; these are well-circumscribed and are measured in millimeters.




Fig. 3. Transverse cross-section computed tomography scan passing through the sacroiliac joints. It shows small, sclerotic, round opacities distributed symmetrically along sacrum, hip bone, and sacroiliac joints.




Fig. 4. Transverse cross-section computed tomography scan passing through the femoral head. It shows numerous hyperdense lesions that conform with the osteopoikilosis in the femoral head; lesions are well-circumscribed and are measured in millimeters.

Authors' Info

Corresponding Author: Giuseppe Guglielmi, MD, Professor; address: Viale L.Pinto 1, 71121 Foggia, Italy; ORCID: 0000-0002-4325-8330; e-mail: giuseppe.guglielmi@unifg.it

Co-authors:

Paparella Maria Teresa, MD; ORCID: 0000-0003-2573-9509; e-mail: mt.paparella@gmail.com

Gangai Ilaria, MD; ORCID: -; e-mail: hilary_ps@libero.it

Porro Chiara, MD; ORCID: 0000-0002-7526-6968; e-mail: chiara.porro@unifg.it

Eusebi Laura, MD; ORCID: 0000-0002-4172-5126; e-mail: lauraeu@virgilio.it

Silveri Ferdinando, MD; ORCID: -; e-mail: fsilveri@libero.it

Cammarota Aldo, MD; ORCID: 0000-0003-4211-5140; e-mail: aldo.cammarota@crob.it

REFERENCES

1. Negi RS, Manchanda KL, Sanga S, et al. Osteopoikilosis – spotted bone disease. *Med J Armed Forces India* 2013;69:196–8.
2. Mahboubia J, Mondher G, Amira M, et al. Osteopoikilosis: a rare cause of bone pain. *Caspian J Intern Med* 2015;6:177–9.
3. Carpintero P, Abad JA, Serrano P, Serrano JA, Rodríguez P, Castro L. Clinical features of ten cases of osteopoikilosis. *ClinRheumatol* 2004; 23: 505–08
4. Tong EC, Samii M, Tchang F. Bone imaging as an aid for the diagnosis of osteopoikilosis. *ClinNuclMed*. 1988;13:816-819.

5. Drouin CA, Grenon H. The association of Buschke-Ollendorfsyndrome and nail-patella syndrome. *J Am Acad Dermatol*. 2002;46:621-625.
6. Albers-Schönberg HE. *Fortschr Roentgen*. 1915;23:174
7. Hellemans J, Preobrazhenska O, Willaert A, et al. Loss-of-function mutations in LEMD3 result in osteopoikilosis, Buschke-Ollendorff syndrome and melorheostosis. *Nat Genet* 2004;36:1213-8.
8. Gutierrez D, Cooper KD, Mitchell AL, et al. Novel somatic mutation in LEMD3 splice site results in Buschke-Ollendorff syndrome with polyostotic melorheostosis and osteopoikilosis. *Pediatr Dermatol* 2015;32:e219-20.
9. Vanhoenacker EM, De Beuckeleer LH, Wan Hul W, Balemans W, Tan GJ, Hill SC, et al. Sclerosing bone dysplasias: genetic and radioclinical features. *Eur Radiol* 2000;10:1423-33.
10. Amezcua-Guerra LM, Mansilla Lory J, Fernandez Tapia S, Pijoan C, Leboeiro I, Martínez-Lavín M, et al. Osteopoikilosis in an ancient skeleton: more than a medical curiosity. *Clin Rheumatol* 2005;24:502-6.
11. Niwayama G, Enostosis, hyperstosis, and periostitis. In: Resnick D, ed. *Diagnosis of Bone and Joint Disorders*. Philadelphia: WB Saunders;1988:4084-4088.
12. Dahan S, Bonafé JL, Laroche M, Rousseau H, Railhac JJ. Iconography of Buschke-Ollendorff syndrome: x ray computed tomography and nuclear magnetic resonance of osteopoikilosis. *Ann Dermatol Venereol*. 1989;116(3):225-230.
13. Mungovan JA, Tung GA, Lambiase RE, Noto RB, Davis RP. Tc-99m MDP uptake in osteopoikilosis. *Clin Nucl Med*. 1994;19(1):6-8.